

cut along dotted line

1

NSMHPCN Pocket Guide COVID-19 EOL DYSPNEA



DOSING CONSIDERATIONS:
Doses are higher dt rapid escalation of COVID.

• **Opioid naïve**

Morphine

Start with 2.5 - 5mg PO or 1-2.5mg subcut/IV q4hrs
PLUS 2.5-5mg PO or 1-2mg subcut/IV q1hr PRN for dyspnea

Hydromorphone

Start with 0.5- 1mg PO or 0.5mg subcut/IV q4hrs PLUS 0.5-1mg PO or 0.5mg subcut/IV q1hr PRN for dyspnea

***Already taking Opioids**

Increase previous opioid by 25%
calculation for BT dose 10% 24 hr total. Give q1h prn if PO, q 30min if Sq

Palliative Nursing Support Line (705) 329-0340 | (844) 429-0340

cut along dotted line

2

NSMHPCN Pocket Guide COVID-19 EOL DYSPNEA



Avoid morphine for moderate/severe renal impairment.
Consider reduced dose (consider by half) in the frail/elderly and those with severe heart lung or neurological diseases.
May consider dosing Q6H instead of Q4H.
MD to review is using 4 or more PRN's.

TITRATE

Increase regularly scheduled dose by 50%, monitor.
Rate is titration depends on how patient tolerating (eg somnolence)
Increase prn by 50% per dose if tolerating opioid well.
Titration can be done after being 18 to 24 hrs ont he same total daily dose.

NAUSEA

Metoclopramide 5-10mg SC/IV/OR mini-bag OR slow infusion QID PRN for opioid induced nausea

Palliative Nursing Support Line (705) 329-0340 | (844) 429-0340

Fold Here

cut along dotted line

3

NSMHPCN Pocket Guide ANXIETY/AGITATION



Associated Anxiety

If opioids not effective or dyspnea crisis:

Methotrimeprazine 6.25mg or 12.5mg
OR

Midazolam 1mg to 2mg SC/IV q1hr PRN

MD to review if 4 or more PRN's in 24 hrs

Palliative Nursing Support Line (705) 329-0340 | (844) 429-0340

4

NSMHPCN Pocket Guide MEDICATIONS



If symptoms Refractory - consider PALLIATIVE SEDATION
(lowest dose of medication to reduce LOC & relieve symptoms), monitor with RASS.

1 ST LINE

Option 1: Methotrimeprazine (Nozinan™)

STAT-25mg subcut (12.5mg in frail, elderly patients).
ATC- 12.5-25mg subcut q4hrs or q6hrs
PRN midazolam 2.5mg or 5mg subcut or IV q30 min
if >4 /24 re-evaluate

Option 2: Midazolam subcut intermittent injections*

STAT-2.5mg or 5mg subcut or IV STAT.
ATC - 2.5mg or 5mg subcut or IV q4hrs.
PRN order of midazolam 1 – 5mg subcut or IV q30 - q 60 min

Palliative Nursing Support Line (705) 329-0340 | (844) 429-0340

Fold Here

cut along dotted line

5

NSMHPCN Pocket Guide MEDICATIONS



Option 3: Midazolam by continuous infusion (C.I.)

STAT - 2.5mg or 5mg subcut or IV stat.
START C.I. at 0.5mg - 1mg/hour subcut or IV by infusion pump. –
Titrate up (or down) q 30 - 60 minutes if needed until the required LOC.

The usual dose required is between 1-5mg/hr. Higher doses may be required in select cases. - If titration required to achieve desired goal (comfort), increase in increments of 0.5mg or 1mg/hr.

If crises occur, may give a bolus doses of midazolam 2.5mg or 5mg subcut or IV q 30 minutes PRN. • If doses of greater than 8-10 mg/hr are required, reassess and consider adding methotrimeprazine

Palliative Nursing Support Line (705) 329-0340 | (844) 429-0340

6

NSMHPCN Pocket Guide MEDICATIONS



2ND LINE

I If options 1, 2, 3 are ineffective
- Add phenobarbital to methotrimeprazine or midazolam.

STAT -60mg, 90mg or 120mg subcut or IV
(depending on the severity of the situation)
-60mg subcut BID.

Long-half life though does not allow for rapid titration (only increase dose every day or 2, not sooner)

Palliative Nursing Support Line (705) 329-0340 | (844) 429-0340

Fold Here

cut along dotted line

cut along dotted line

cut along dotted line

cut along dotted line

cut along dotted line

cut along dotted line

cut along dotted line

7

NSMHPCN Pocket Guide

RICHMOND AGITATION AND SEDATION - RASS



Step	Procedure	Score
1	Observe Patient - Patient is alert, restless or agitated	0 to +4
2	If not alert, state patient's name and say to open eyes and look at speaker. - Patient awakens with sustained eye opening and eye contact. - Patient awakens with eye opening and eye contact but not sustained. - Patient has any movement in response to voice but no eye contact	-1 -2 -3
3	If patient does not respond to voice, physically stimulate patient by shaking shoulder and/or rubbing sternum*: - Patient has any movement to physical stimulation. - Patient has no response to any stimulation.	-4 -5

*Rubbing the sternum is not appropriate for palliative care patient assessment and is not recommended.

Palliative Nursing Support Line (705) 329-0340 | (844) 429-0340

cut along dotted line

8

NSMHPCN Pocket Guide

RICHMOND AGITATION AND SEDATION - RASS



Scale	Label	Description
+4	Combative	Violent, immediate danger to staff
+3	Very agitated	Pulls or removes tube(s) or catheter(s); aggressive
+2	Agitated	Frequent non-purposeful movement, fights ventilator
+1	Restless	Anxious but movements not aggressive, vigorous
0	Alert and calm	Spontaneously pays attention to care giver
-1	Drowsy	Not fully alert, but has sustained awakening (eye-opening/eye contact) to voice (>10 seconds)
-2	Light sedation	Briefly awakens with eye contact to voice (<10 seconds)
-3	Moderate sedation	Movement or eye opening to voice (but no eye contact)
-4	Deep sedation	No response to voice, but movement or eye opening to physical stimulation
-5	Unarousable	No response to voice or physical stimulation

Palliative Nursing Support Line (705) 329-0340 | (844) 429-0340

cut along dotted line

cut along dotted line

Fold Here

9

NSMHPCN Pocket Guide

SELF-CARE



Practicing self-care is important to ease the physical affects of being under added stress of the COVID-19 pandemic. A few ways to observe self-care are:

- Following a routine - Putting a new routine in place is an important part of ensuring one is able to adjust well to having the old routine changed. This includes waking up at a normal time, getting dressed, brushing teeth/hair, etc.
- Proper Nutrition - Continuing to follow a balanced diet will help ensure that the body is receiving the nutrients that it needs in order to maintain a healthy immune system.
- Sleep - Following a sleep routine, in conjunction with proper nutrition, is important to ensure one is able to continue to be as comfortable as possible.

Palliative Nursing Support Line (705) 329-0340 | (844) 429-0340

10

NSMHPCN Pocket Guide

SELF-CARE



- Physical Activity - Many may find that the options for physical activity have become minimal. There are options that individuals can do in the comfort of their own home/neighbourhood.
 - Yoga - there are videos available on YouTube for all levels of yoga. Apps such as Down Dog are available for download on all operating systems.
 - Walking trails - the majority of walking trails remain open for use as long as physical distancing is being observed.
 - Stairs - for those who are unable to leave their homes there is the option of doing reps of going up and down the stairs.

Self-Care Resources:

- Down Dog App
- Calm Meditation App
- Bounce Back Ontario
- Regional Bereavement Support Line (705) 325-7871
- Kids Help Phone 1-800-668-6868
- Indigenous Hope for Wellness: 1-855-242-3310

Palliative Nursing Support Line (705) 329-0340 | (844) 429-0340

cut along dotted line

cut along dotted line

Fold Here

cut along dotted line

cut along dotted line